
## 2016 Disease Rating\* Rhizomania



<sup>\*</sup> The disease ratings are general in nature and may not be representative of all fields in a township

Moderate Severe



## Rhizomania Management\*



Disease Severity	Management Opportunities
Severe	<ul> <li>Plant only Dual, Tandem, or Stacked resistant varieties</li> <li>Use good drainage practices</li> <li>Plant as early as possible</li> <li>Maximize rotation length</li> <li>Practice proper crop rotation</li> <li>Monitor for blinkers and resistance breaking strains</li> <li>Watch for severe sugar content decline</li> <li>Tile drainage very likely beneficial</li> <li>Consider renting land with less severity</li> </ul>
Moderate	<ul> <li>Plant only Dual, Tandem, or Stacked resistant varieties</li> <li>Use good drainage practices</li> <li>Plant as early as possible</li> <li>Maximize rotation length</li> <li>Practice proper crop rotation</li> <li>Monitor for blinkers and resistance breaking strains</li> <li>Watch for sugar content decline</li> <li>Monitor field symptoms for increased severity</li> </ul>
Slight	<ul> <li>Plant any resistant variety</li> <li>Use good drainage practices</li> <li>Plant as early as possible</li> <li>Practice proper crop rotation</li> <li>Monitor field symptoms for increased severity</li> <li>Watch for sugar content decline</li> </ul>
	<ul> <li>Prevent disease introduction</li> </ul>

\* Management strategies listed in priority order

None

Use good drainage practicesMonitor field symptoms

Practice proper crop rotation